## antiviral and associated therapy - all results

Treatment	Number of studies	Demonstrated or suggested results	Inconclusive results	Uncertain results	Safety results
chloroquine and derivatives	4 studies <sup>1</sup>		inconclusive results for : death D28; deaths; deaths (time to event analysis only); clinical deterioration; clinical improvement (time to event analysis only); hospital discharge; ICU admission; off oxygenation	suggested 86	
ivermectin	3 studies <sup>2</sup>		inconclusive results for : deaths; deaths (time to event analysis only); clinical improvement; clinical improvement (7-day); viral clearance; ICU admission	suggested 86	
lopinavir/ritonavir	2 studies <sup>3</sup>		inconclusive results for : deaths; deaths (time to event analysis only); clinical improvement; clinical improvement (14-day); clinical improvement (28-day); clinical improvement (7-day); clinical improvement (time to event analysis only); death or ventilation; hospital discharge; serious adverse events		
azithromycin	1 study <sup>4</sup>		inconclusive results for : death D28; deaths; deaths (time to event analysis only); clinical improvement; clinical improvement (14-day); clinical improvement (28-day); clinical improvement (7-day); related SAE (TRSAE); serious adverse events; long QT; renal impairment		
favipiravir	1 study <sup>5</sup>		inconclusive results for : deaths; hospital discharge		

Continued on next page

Treatment	Number of studies	Demonstrated or suggested results	Inconclusive results	Uncertain results	Safety results
Lopinavir/ritonavir plus hydroxychloroquine	1 study <sup>6</sup>		inconclusive results for : clinical improvement ; clinical improvement (14-day); hospital discharge; serious adverse events	suggested 42	
remdesivir	1 study <sup>7</sup>		inconclusive results for : deaths; clinical deterioration; clinical improvement; clinical improvement (14-day); clinical improvement (28-day); clinical improvement (7-day); clinical improvement (time to event analysis only); AE leading to drug discontinuation; serious adverse events; deep vein thrombosis; elevated liver enzymes; hyperbilirubinemia; pulmonary embolism; renal impairment		
sofosbuvir and daclatasvir	1 study <sup>8</sup>		inconclusive results for : adverse events	suggested 88	

Uncertain results are statistically significant results but obtain in trial with high risk of bias.

Demonstrated results are significant results obtained on the primary endpoint of a trial at low risk of bias or with some concerns.

## **Notes**

<sup>&</sup>lt;sup>1</sup>HYDRA (Hernandez-Cardenas), 2021 (NCT04315896); REMAP-CAP-HCQ, 2020 (NCT02735707); Mahevas, 2020 (); Yu, 2020 ()

<sup>&</sup>lt;sup>2</sup>Galan, 2021 (RBR-8h7q82); Okumuş, 2020 (NCT04646109); Camprubi, 2020 ()
<sup>3</sup>Cao, 2020 (ChiCTR2000029308); REMAP-CAP (lopinavir/ritonavir only), 2020 (NCT02735707)

<sup>&</sup>lt;sup>4</sup>COALITION II Covid-19 Brazil (Furtado), 2020 (NCT04321278)

<sup>&</sup>lt;sup>5</sup>Kocayigit H, 2020 ()

<sup>&</sup>lt;sup>6</sup>REMAP-CAP (lopinavir/ritonavir plus hydroxychloroquine), 2020 (NCT02735707)

<sup>&</sup>lt;sup>7</sup>CAP-China (Wang et al.), 2020 (NCT04257656)

<sup>&</sup>lt;sup>8</sup>Eslami, 2020 (IRCT20200324046850N2)